

Diabrotica speciosa

Scientific name

Diabrotica speciosa Germar

Synonyms:

Diabrotica amabilis, *Diabrotica hexaspilota*, *Diabrotica simoni*, *Diabrotica simulans*, *Diabrotica vicens*, and *Galeruca speciosa*

Common names

Cucurbit beetle, chrysanthemum beetle, San Antonio beetle, and South American corn rootworm

Type of pest

Beetle

Taxonomic position

Class: Insecta, **Order:** Coleoptera, **Family:** Chrysomelidae

Reason for Inclusion in Manual

CAPS Target: AHP Prioritized Pest List - 2010

Pest Description

Diabrotica speciosa was first described by Germar in 1824, as *Galeruca speciosa*. Two subspecies have been described, *D. speciosa vicens* (Bolivia, Peru and Ecuador), and *D. speciosa amabilis* (Bolivia, Colombia, Venezuela and Panama). These two subspecies differ mainly in the coloring of the head and elytra (Araujo Marques, 1941; Bechyne and Bechyne, 1962).

Eggs: Eggs are ovoid, about 0.74 x 0.36 mm, clear white to pale yellow. They exhibit fine reticulation that under the microscope appears like a pattern of polygonal ridges that enclose a variable number of pits (12 to 30) (Krysan, 1986). Eggs are laid in the soil near the base of a host plant in clusters, lightly agglutinated by a colorless secretion. The mandibles and anal plate of the developing larvae can be seen in mature eggs.

Larvae: Defago (1991) published a detailed description of the third instar of *D. speciosa*. First instars are about 1.2 mm long, and mature third instars are about 8.5 mm long. They are subcylindrical; chalky white; head capsule dirty yellow to light brown, epicraneal and frontal sutures lighter, with long light-brown setae; mandibles reddish dark brown; antennae and palpi pale yellow. Body covered by sparse, short, dark setae; light brown irregular prothoracic plate; dark brown anal plate on the ninth segment, with a pair of small urogomphi. A pygopod is formed by the tenth segment, which serves as a locomotion and adherence organ.

Pupae: Pupae are 5.8 to 7.1 mm long and white. Females with a pair of tubercles near the apex. Mature third instars build an 8 x 4 mm oval cell in the soil in which they pupate, and teneral remain for about 3 days.

Adults: Full descriptions of *D. speciosa* are given by Baly (1886), Araujo Marques (1941), and Christensen (1943). Adults are 5.5 to 7.3 mm long; antennae 4 to 5 mm (Fig. 1). General color grass-green (USDA, 1957); antennae filiform and dark (reddish-brown to black) and nearly equal to the body in length, first three basal segments lighter; head ranging from reddish brown to black; labrum, scutellum, metathorax, tibiae and tarsi black; elytra each with three large oval transverse spots, basal spots larger and usually reddish toward the humeral callus, the rest yellow.



Figure 1. Adult *Diabrotica speciosa*. Photo courtesy of Hernan Tolosa.

Ventrally, head and metathorax dark brown, prothorax green, mesothorax and abdomen light brown or yellow-green. Pronotum bi-foveate, convex, smooth, shiny, $\frac{1}{4}$ wider than long. Male antennae proportionally longer than female antennae. Males with an extra sclerite on the apex of the abdomen that makes it look blunt, compared with the rather pointed female apex.

Biology and Ecology

Eggs are laid on the soil near a larval host plant. An approximately 92% success rate at 27°C is takes place after about 8 days. *Diabrotica speciosa* undergoes three larval instars, which are easily differentiated by the size of the head capsule (see larval description above). In laboratory tests, maize was included in the grouping of most suitable hosts (along with wheat and peanuts), in terms of survival from egg to adult (Cabrera Walsh, 2003). First instars are normally scattered throughout the host's root system, but mature larvae tend to congregate in the upper 10 cm of the root under the crown. The larval stage lasts 23 to 25 days (~12 days in laboratory conditions at 25°C), including an inactive prepupal period of 2 to 3 days. At 25°C, the pupal stage lasts 6 days, and is followed by a period of 3 to 5 days during which the recently molted adults remain in the pupal cell, presumably for the cuticle to tan (USDA, 1957).

Young beetles have a yellowish or pale brown color, which turns green with bright yellow spots in 3 days if fresh food is provided. Under laboratory conditions, mating has been observed between 4 and 6 days after emergence, and some females were observed mating again at day 35. Each female laid an average of 1164 eggs during her lifetime, starting on day 8 and extending for a maximum of 77 days. Peak oviposition was observed on days 16 through 56. In a laboratory environment, oviposition on

maize was preferred over pumpkin, potato and bean seedlings, and maize was as attractive as peanuts in choice tests (Cabrera Walsh, 2003). The number of overlapping generations is conditioned by latitude and climate, being continuous in tropical areas. In Buenos Aires, Argentina, observations indicate there are about three generations per year; the number and timing depends on latitude and climate. Overwintering occurs as an adult (USDA, 1957). These adults can be found concealed in the rosette and crown of winter-growing plants, and they are fairly cold-tolerant (EPPO, 2005).

Pest Importance

Diabrotica speciosa is considered to be an important pest throughout southern South America (except Chile), but, being highly polyphagous, qualitative reports of its impact on different crops vary in different regions. It is considered an important pest of maize, cucurbits, and orchard crops throughout its distribution (CABI, 2007). Although it migrates as an adult, no information on observed distances has been found. Redistributing soil via farm machinery that is contaminated with eggs and-or pupae is also a concern.

Adults of this chrysomelid feed on foliage, pollen, flowers and fruits of many plants. The larvae are pests of roots, especially maize. It is the most harmful species of *Diabrotica* in Argentina, mainly affecting peanuts in the center of the country. It causes considerable damage to watermelon, squash and tomatoes in Brazil, and potatoes and wheat in southeast Brazil. Young squash plantings and immature tomato fruits are severely damaged in Brazil. Populations are so heavy in some years in Paraguay that vegetable crops are almost completely destroyed. Severe injury also occurs on flowers of various ornamentals such as dahlias and chrysanthemums (USDA, 1957). Economic thresholds of two insects per plant for *Phaseolus vulgaris* were determined by Pereira et al. (1997).

IPM programs to combat *D. speciosa* in South America recommend no-till cultural practices, insecticides when reaching economically damaging levels and a rotation of maize, wheat, and soybeans. In South America, insecticides (carbamates, organophosphates and, more recently, tefluthrin and chlorethoxyfos) to control larvae and baits (along with broad-spectrum insecticides) to control adults are widely used. These baits are sliced roots of several different wild cucurbits laced with insecticides.

Although there is research into using parasitoids (brachonids and tachinids) and pathogens (*Beauveria* spp. and *Metarhizium anisopliae*) to combat this pest, no successful biological control programs have been mentioned.

Symptoms/Signs

The larval damage resulting from root feeding can cause host death when the host is small, but the larvae will usually only induce stunted growth in larger host plants, due to a reduction in nutrient uptake. Wheat is a larval host and stunting would be the primary symptom. In corn, attack on young plants by larvae produces a typical condition known as 'goose neck', in which the plant exhibits stunted growth, reduced vigor, and the first few internodes of the plant grow bent, sometimes to such an extent that the plant

actually lies on the ground (Figure 2). In the case of peanuts and potatoes, the larvae cause external damage or short bores, similar to those of several other pests such as wireworms and other chrysomelids.

On corn, the most economically important stage is the adult, which feeds on the tassels, preventing pollination and kernel number. Adults also cause defoliation and general feeding damage to leaves, flowers and fruit

(EPPO, 2005). Like other *Diabrotica* spp., they are especially associated with Cucurbitaceae and are tolerant of cucubitacins and generally feed on pollen-rich plant structures of over 70 plant species. When flowers are scarce, beetles may feed on the tender green parts of other hosts, such as alfalfa, potatoes, corn, bean, soybean, lettuce, and cabbage (EPPO, 2005).



Figure 2. 'Gooseneck' growth form of corn. Photo courtesy of The Ohio State University.



Figure 3. Grape cluster after a severe outbreak of *D. speciosa* during the bloom period (A) and normal cluster (B). Photos courtesy of Roberto et al. (2001).

In grape, adult beetles eat young leaf edges during budding, which usually does not seriously damage the host (Roberto et al., 2001). During the blooming period, however, beetles have been observed on flowers eating the style, stigma, and eventually the ovary. Beetle stigma feeding determines flower aborting and, as a consequence,

clusters show low numbers of flowers and fruits (Fig. 3). Weedy hosts need to be controlled as beetles can also be observed feeding on and moving into grape from surrounding weeds.

Known Hosts

Root-feeding larvae of *D. speciosa* are polyphagous, but the known host range includes corn, wheat, peanut, soybean, and potato. Cabrera Walsh (2003) found that larvae developed well on corn, peanut, and soybean roots, but not so well on pumpkin, beans, and potato. Oviposition preferences roughly parallel larval suitability, but there was a clear preference for cucurbits as adult food, when available; pigweed, sunflower, and alfalfa are secondary hosts. As an adult, *D. speciosa* has been reported feeding on more than 70 host species (Christensen, 1943; Heineck-Leon and Salles, 1997).

Major hosts

Arachis spp. (peanut), *Capsicum* spp. (pepper), *Cucurbita maxima* (winter squash), *Cucurbita pepo* (ornamental gourd), *Glycine max* (soybean), *Solanum tuberosum* (potato), *Triticum* spp. (wheat), *Vitis vinifera* (grape), and *Zea* spp. (corn).

Minor hosts

Allium spp. (onion, leek), *Alternanthera philoxeroides* (alligatorweed), *Amaranthus* spp. (pigweeds), *Apium graveolens* (celery), *Artemisia* spp. (absinthium, tarragon), *Asparagus* spp. (asparagus), *Avena* spp. (oats), *Baccharis articulata*, *Beta vulgaris* (beet), Brassicaceae (mustards), *Bromus catharticus* (prairie grass), *Carica papaya* (papaya), *Cayaponia* spp., *Chenopodium* spp., *Chrysanthemum* spp., *Cichorium* spp. (chicory, endive), *Citrullus vulgaris* (watermelon), *Citrus* spp., *Coriandrum sativum* (coriander), *Coronopus didymus* (twin cress), *Cucumis* spp. (melons, cucumbers, gerkins), Cucurbitaceae (cucurbits), *Cucurbitella asperata*, *Cynara* spp. (artichoke), *Cynodon dactylon* (Bahama grass), *Cyphomandra betacea* (tree tomato), *Dahlia pinnata* (pinnate dahlia), *Datura* spp., *Daucus carota* (carrot), *Fragaria vesca* (wild strawberry), *Gossypium* spp. (cotton), *Helianthus annuus* (sunflower), *Helianthus tuberosus* (Jerusalem artichoke), *Hibiscus* spp., *Ilex paraguayensis* (Paraguay tea), *Ipomoea* spp. (sweet potato, morning glory), *Lactuca sativa* (lettuce), *Lagenaria siceraria* (bottle gourd), *Lavandula officinalis* (English lavender), *Lilium maculatum* (sukash-yuri), *Linum usitatissimum* (flax), *Lolium perenne* (rye grass), *Luffa* spp. (loofah), *Lycopersicon esculentum* (tomato), *Malus* spp. (apple), *Malva* spp. (mallow), *Matricaria chamomilla* (chamomile), *Medicago sativa* (alfalfa), *Melilotus albus* (yellow sweet clover), *Mentha* spp. (mint), *Morrenia odorata* (latexplant), *Musa* spp. (banana), *Nasturtium officinale* (watercress), *Nicotiana tabacum* (tobacco), *Ocimum basilicum* (basil), *Origanum vulgare* (oregano), *Oryza sativa* (rice), *Passiflora coerulea* (passion flower), *Petroselinum crispum* (parsley), *Pharbitis purpurea* (common morning glory), *Phaseolus* spp. (beans), *Physalis viscosa* (starhair groundcherry), *Pimpinella anisum* (anise), *Pisum sativum* (pea), *Prunus* spp. (stone fruit), *Raphanus sativus* (radish), *Rosa* spp. (rose), *Sechium edule* (chayote), *Sicyos polycanthus*, *Solanum* spp., *Solidago chilensis* (goldenrod), *Sorghum* spp., *Spinacia oleracea* (spinach), *Taraxicum officinale* (dandelion), *Thea sinensis* (tea), *Trifolium* spp. (clover), *Triticum* spp., *Tropaeolum majus* (Nasturtium), and *Zingiber officinale* (ginger).

Known Vectors (or associated organisms)

There is evidence that *D. speciosa* is a viral vector for comoviruses, southern bean mosaic virus, mimosa mosaic virus, tymoviruses (such as passionfruit yellow mosaic virus), carmoviruses, and purple granadilla mosaic virus (Ribeiro et al., 1996; Germain, 2000). Lin et al. (1984) showed that *D. speciosa* transmitted cowpea severe mosaic virus (CPSMV – comovirus) to bean. Ribeiro et al. (1996) showed that eggplant mosaic virus (EMV – tymovirus) was transmitted to tobacco by *D. speciosa*. Cabrera Walsh (2003) mention that *D. speciosa* may also transmit bacterial wilt, caused by *Erwinia tracheiphila*, in cucurbits.



Figure 4. Adult banded cucumber beetle, *Diabrotica balteata*. Photo courtesy of John L. Capinera, University of Florida

Known Distribution

Central America: Costa Rica and Panama. **South America:** Argentina, Bolivia, Brazil, Columbia, Ecuador, French Guiana, Paraguay, Peru, Uruguay, and Venezuela. There is a record of *D. speciosa* from Mexico, but according to Krysan (1986), it is almost certainly an error.

Potential Distribution within the United States

As of 2004, *Diabrotica speciosa* has been intercepted over 300 times at ports of entry in the United States, but little is known on its potential distribution within the United States. According to a recent host analysis by USDA-APHIS-PPQ-CPHST, the greatest risk for establishment of *D. speciosa* based solely on the presence of hosts occurs in portions of Arkansas, Colorado, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Mississippi, Missouri, Nebraska, North Carolina, North Dakota, Ohio, Oklahoma, Texas, and Wisconsin. The pest occurs from temperate Argentina to tropical Brazil. The polyphagous nature of *D. speciosa* increases the likelihood of finding hosts and suitable environment if it were introduced into the United States, and is thought to be able to adapt to more temperate climates.

Survey

CAPS-Approved Method: Visual.

Literature-Based Methods: Visual survey: Visual detection of adults is easy, as their feeding period spans from dawn until dusk. Detection of larval damage, on the other hand, is more difficult. First instars are very difficult to sample, and even large infestations can go undetected until the damage caused to the host is extensive. Larger larvae can sometimes be observed feeding on the roots of plants immediately after pulling out of the soil, but methodical sampling and counting methods have not been developed, as they have been for the North American pest species (Fisher and Bergman, 1986).

Trapping: Adults *D. speciosa* appear to be universally attracted to aromatic compounds from squash blossoms, though the specific compound(s) that attract the beetles varies from species to species. Often, simple blends of two or three compounds are much more potent attractants than any single compound. In addition, female-produced sex attractant pheromones are used for mate location in this genus. In a preliminary trapping test in Brazil, a number of squash volatiles were screened for potential attraction, and 1,4-dimethoxybenzene showed promise as an attractant for *D. speciosa* (Ventura et al., 2000). Traps baited with 1,4-dimethoxybenzene, a volatile substance of *Cucurbita maxima* blossoms captured 29.4 times and 9.4 times more beetles than controls in soybean and common bean fields, respectively (Ventura et al., 2000).



Figure 5. Western corn rootworm, *Diabrotica virgifera*. Courtesy of USDA-ARS.



Figure 6. Southern corn rootworm, *Diabrotica undecimpunctata*. Courtesy of Clemson University - USDA Cooperative Extension Slide Series, www.bugwood.org.

The USDA-CPHST laboratory in Otis, MA has applied for funding to manufacture and test potential lures for *D. speciosa*, but has yet to begin work toward this goal.

Key Diagnostics/Identification

CAPS-Approved Method: Confirmation of *D. speciosa* is by morphological identification. *Diabrotica speciosa* is almost identical to *D. balteata* (Fig. 4), which is widely present in the southern United States. Confirmation by a chrysomelid specialist is required. *D. speciosa* can also be confused with *Diabrotica viridula* (not present in the United States) and other pestiferous *Diabrotica* species in South America.

Literature-Based Methods:

Diabrotica speciosa somewhat resembles the other main pestiferous *Diabrotica* in South America, *D. viridula*, in coloring, size, biology and host range; but *D. viridula* has dark brown areas toward the cephalic edge of the elytral spots, and distinct humeral plicae. Also, the larvae of *D. viridula* lack urogomphi on the anal plate.

Easily Confused Pests

Survey and detection based on visual detection of symptoms is quite difficult and many other pests can be easily confused. Symptoms, such as dead heart in wheat, goose neck in maize, or stunted growth in most of the larval hosts of *D. speciosa*, could be attributed to several other root feeders, such as wireworms (*Conoderus* spp.; Elateridae), white grubs, (*Phytalus* spp., *Cyclocephala* spp., *Diloboderus abderus*; Melolonthidae), *Pantomorus* spp. and *Listronotus bonariensis* (Curculionidae), and several chrysomelids (*Caeporis* spp., *Colaspis* spp., *Maecolaspis* spp., *Diphaulaca* spp. and *Cerotoma arcuata*) (Gassen, 1984, 1989).

Other rootworms (western corn rootworm, southern corn rootworm) are easily distinguished from *D. speciosa* as adults by the markings on elytra (compare Figs. 1, 5 and 6).

References

- Araujo Marques, M.** 1941. Contributio ao estudo dos crisomeledeos do gunero *Diabrotica*. Bol. Escola Nac. Agron., 2:61-143. (In Portuguese)
- Baly, J.S.** 1886. The Colombian species of the genus *Diabrotica*, with descriptions of those hitherto uncharacterized. Part I. Zoological Journal of the Linnean Society, 19:213-229.
- Bechyne, J. and Bechyne, B.** 1962. Liste der bisher in Rio Grande do Sul gefundenen Galeruciden. Pesquisas (Zool.), 6:1-63.
- CABI.** 2007. Crop Protection Compendium. Wallingford, UK: CAB International. www.cabicompndium.org/cpc.
- Cabrera Walsh, G.** 2003. Host range and reproductive traits of *Diabrotica speciosa* (Germar) and *Diabrotica viridula* (F.) (Coleoptera: Chrysomelidae) two species of South American pest rootworms, with notes on other species of Diabroticina. Environmental Entomology 32(2): 276-285.
- Christensen, J.R.** 1943. Estudio sobre el g'nero Diabrotica Chev. en la Argentina. Rev. Facultad de Agronomia y Veterinaria, 10:464-516. (In Spanish)
- Defago, M.T.** 1991. Caracterizacion del tercer estadio larval de *Diabrotica speciosa*. Rev. Peruana de Ent., 33:102-104. (In Spanish)

EPPO. 2005. *Diabrotica speciosa*. OEPP/EPPO Bulletin 35: 374-376.

Fisher, J.R. and Bergman, M.K. 1986. In: [Krysan JL, Miller TA, eds.] Methods for the Study of Pest *Diabrotica*. New York, USA: Springer.

Gassen, D.N. 1984. Insetos associados a cultura do trigo no Brasil. Circular Tecnica, 3. Passo Fundo, RS, Brazil: EMRAPA-CPNT.

Gassen, D.N. 1989. Insetos subterraneos prejudiciais as culturas no sul do Brasil. EMBRAPA-CNPT. Documentos, 13. Passo Fundo, RS, Brazil: EMBRAPA-CNPT.

Germain, J.F. 2000. *Diabrotica speciosa* (Germar), PRA Area: Europe (01/8504). 1-3. European and Mediterranean Plant Protection Organization.

Heineck-Leonel, M.A. and Salles, L.A.B. 1997. Incidence of parasitoids and pathogens in adults of *Diabrotica speciosa* (Germ.) (Coleoptera: Chrysomelidae) in Pelotas, RS. Anais da Sociedade Entomológica do Brasil, 26(1):81-85.

Krysan, J.L. 1986. Introduction: biology, distribution, and identification of pest *Diabrotica*. In: [Krysan JL, Miller TA, eds.] Methods for the Study of Pest *Diabrotica*. New York, USA: Springer.

Lin, M.T., Hill, J.H., Kitajima, E.W., and Costa, CL. 1984. Two new serotypes of cowpea severe mosaic virus. Phytopathology 74: 581-585.

Pereira, M.F.A., Delfini, L.G., Antoniacomi, M.R., and Calafiori, M.H. 1997. Damage caused by the leaf beetle, *Diabrotica speciosa* (Germar, 1824), on beans (*Phaseolus vulgaris* L.), with integrated management. Ecosystema, 22:17-20.

Ribeiro, S.G., Kitajima, E.W., Oliveira, C.R.B., and Koenig, R. 1996. A strain of eggplant mosaic virus isolated from naturally infected tobacco plants in Brazil. Plant Dis. 80: 446-449.

Roberto, S.R., Genta, W., and Ventura, M.U. 2001. *Diabrotica speciosa* (Ger.) (Coleoptera: Chrysomelida): New Pest in Table Grape Orchards. Neotropical Entomology 30(4): 721-722.

USDA, 1957. Cooperative Economic Insect Report, 7(2):5-6.

Ventura, M.U., Martins, M.C., and Pasini, A. 2000. Response of *Diabrotica speciosa* and *Cerotoma acuata tingomariana* (Coleoptera: Chrysomelidae) to volatile attractants. Florida Entomologist 83(4): 403-410.